arXiv:nlin/0302026v1 [nlin.SI] 12 Feb 2003

Classical transcendental solutions of the Painlevé equations and their degeneration

Tetsu Masuda
Department of Mathematics, Kobe University,
Rokko, Kobe, 657-8501, Japan
masuda@math.kobe-u.ac.jp

Abstract

We present a determinant expression for a family of classical transcendental solutions of the Painlevé V and the Painlevé VI equation. Degeneration of these solutions along the process of coalescence for the Painlevé equations is discussed.

1 Introduction

It is known that each of the Painlevé equations (except for $\mathrm{P_I}$) admit two classes of classical solutions. One is classical transcendental solutions expressible in terms of special functions of hypergeometric type. Another one is algebraic or rational solutions. It is also known that such classical solutions are located on special places from a viewpoint of symmetry of the affine Weyl group as a Bäcklund transformations group of each Painlevé equation. The rough picture is that classical transcendental solutions exist on the reflection hyperplanes of the affine Weyl group and algebraic (or rational) solutions do on the fixed points with respect to the Bäcklund transformations corresponding to automorphisms of the Dynkin diagram.

In this paper, we concentrate our attention on the classical transcendental solutions. One of the important features of these solutions is that they can be expressed in terms of 2-directional Wronskians or Casorati determinants whose entries are given by the corresponding special function of hypergeometric type. In fact, such determinant expressions for $\mathrm{P_{II}}$, $\mathrm{P_{III}}$ and $\mathrm{P_{IV}}$ have been presented [12, 13, 5, 6].

The aim of this paper is to present a determinant expression for a family of classical transcendental solutions of the Painlevé V equation

$$\frac{d^2q}{dt^2} = \left(\frac{1}{2q} + \frac{1}{q-1}\right)\left(\frac{dq}{dt}\right)^2 - \frac{1}{t}\frac{dq}{dt} + \frac{(q-1)^2}{2t^2}\left(\kappa_\infty^2 q - \frac{\kappa_0^2}{q}\right) - (\theta+1)\frac{q}{t} - \frac{q(q+1)}{2(q-1)}, \tag{1.1}$$

which is equivalent to the Hamiltonian system

$$\mathrm{S_V}: \qquad q' = \frac{\partial H}{\partial p}, \quad p' = -\frac{\partial H}{\partial q}, \quad {}' = t\frac{d}{dt}, \tag{1.2}$$

with the Hamiltonian

$$\begin{gathered} H = q(q-1)^2p^2 - \left[\kappa_0(q-1)^2 + \theta q(q-1) + tq\right]p + \kappa(q-1), \\ \kappa = \tfrac{1}{4}(\kappa_0+\theta)^2 - \tfrac{1}{4}\kappa_\infty^2, \end{gathered} \tag{1.3}$$

and the Painlevé VI equation

$$\begin{aligned}\frac{d^2q}{dt^2} = \frac{1}{2}\left(\frac{1}{q}+\frac{1}{q-1}+\frac{1}{q-t}\right)\left(\frac{dq}{dt}\right)^2 - \left(\frac{1}{t}+\frac{1}{t-1}+\frac{1}{q-t}\right)\frac{dq}{dt} \\ +\frac{q(q-1)(q-t)}{2t^2(t-1)^2}\left[\kappa_\infty^2 - \kappa_0^2\frac{t}{q^2}+\kappa_1^2\frac{t-1}{(q-1)^2}+(1-\theta^2)\frac{t(t-1)}{(q-t)^2}\right],\end{aligned} \tag{1.4}$$

which is equivalent to the Hamiltonian system

$$\mathrm{S_{VI}}: \qquad q' = \frac{\partial H}{\partial p}, \quad p' = -\frac{\partial H}{\partial q}, \quad ' = t(t-1)\frac{d}{dt}, \tag{1.5}$$

with the Hamiltonian

$$\begin{aligned} H = q(q-1)(q-t)p^2 - [\kappa_0(q-1)(q-t)+\kappa_1 q(q-t)+(\theta-1)q(q-1)]p + \kappa(q-t), \\ \kappa = \tfrac{1}{4}(\kappa_0+\kappa_1+\theta-1)^2 - \tfrac{1}{4}\kappa_\infty^2,\end{aligned} \tag{1.6}$$

respectively.

Let us explain how to construct a family of classical transcendental solutions of the Painlevé equations. As an example, we take $\mathrm{P_{II}}$:

$$\frac{d^2q}{dt^2} = 2q^3 - 2tq + 2\left(\alpha+\frac{1}{2}\right), \tag{1.7}$$

which is equivalent to the Hamiltonian system

$$\mathrm{S_{II}}: \qquad q' = \frac{\partial H}{\partial p}, \quad p' = -\frac{\partial H}{\partial q}, \quad ' = \frac{d}{dt}, \tag{1.8}$$

with the Hamiltonian

$$H = -p^2 - (q^2-t)p + \alpha q. \tag{1.9}$$

If $\alpha = 0$, the right-hand side of the second equation of (1.8) is divisible by p. This means that $\mathrm{S_{II}}$ admits the specialization of $p = 0$ if $\alpha = 0$. The first equation of (1.8) yields the Riccati equation $q' = -q^2 + t$. Setting $q = (\log\varphi)'$, we get a linear equation $\varphi'' = t\varphi$, which is nothing but Airy's differential equation. Thus, we find that $\mathrm{P_{II}}$ admits, when $\alpha = 0$, the 1-parameter family of special solutions expressed by a rational function of the Airy function and its derivative.

Let introduce the τ-function via the Hamiltonian (1.9) as $H(\alpha) = (\log\tau(\alpha))'$. Then, it is known that a sequence of the τ-functions $\tau_n = \tau(\alpha+n)$ $(n \in \mathbb{Z})$ satisfies the Toda equation

$$\tau_{n+1}\tau_{n-1} = \tau_n''\tau_n - (\tau_n')^2, \tag{1.10}$$

which corresponds to the Bäcklund transformation

$$\alpha \to \alpha - 1, \quad q \to -q - \frac{2\alpha}{q'+q^2-t}. \tag{1.11}$$

We iterate the Bäcklund transformations to the above Riccati solution to obtain a family of classical transcendental solutions. What we have to do is reduced to solving the Toda equation (1.10) with the initial conditions

$$\tau_{-1} = 0, \quad \tau_0 = 1, \quad \tau_1 = \varphi. \tag{1.12}$$

Therefore, we have the following proposition [12].

Proposition 1.1 *(Okamoto) We define the functions* τ_n $(n \in \mathbb{Z}_{\geq 0})$ *by*

$$\tau_n = \begin{vmatrix} \varphi^{(0)} & \varphi^{(1)} & \cdots & \varphi^{(n-1)} \\ \varphi^{(1)} & \varphi^{(2)} & \cdots & \varphi^{(n)} \\ \vdots & \vdots & \ddots & \vdots \\ \varphi^{(n-1)} & \varphi^{(n)} & \cdots & \varphi^{(2n-2)} \end{vmatrix}, \quad \varphi^{(k)} = \left(\frac{d}{dt}\right)^k \varphi, \tag{1.13}$$

where φ *is the general solution of Airy's differential equation* $\varphi'' = t\varphi$*. Then,*

$$q = \frac{d}{dt} \log \frac{\tau_{n+1}}{\tau_n}, \quad p = \frac{\tau_{n+1}\tau_{n-1}}{\tau_n^2}, \tag{1.14}$$

$$\alpha = n, \tag{1.15}$$

gives a family of classical transcendental solutions of S_{II}*.*

By the similar procedures, it is possible to obtain a determinant expression for a family of classical transcendental solutions to $\mathrm{P_V}$ and $\mathrm{P_{VI}}$, which are presented in Section 2 and 3, respectively. As is well known, $\mathrm{P_{VI}}$ degenerates $\mathrm{P_V}, \ldots, \mathrm{P_I}$ by successive limiting procedures [14, 2]. In section 4, we discuss the degeneration of the family of classical transcendental solutions.

2 Classical transcendental solutions of the Painlevé V equation

2.1 The symmetric form of the Painlevé V equation

First, we summarize the symmetric form of the Painlevé V equation [7, 8]. The symmetric form of $\mathrm{P_V}$ is given by

$$\begin{aligned} f_0' &= f_0 f_2 (f_1 - f_3) + \left(\frac{1}{2} - \alpha_2\right) f_0 + \alpha_0 f_2, \\ f_1' &= f_1 f_3 (f_2 - f_0) + \left(\frac{1}{2} - \alpha_3\right) f_1 + \alpha_1 f_3, \\ f_2' &= f_2 f_0 (f_3 - f_1) + \left(\frac{1}{2} - \alpha_0\right) f_2 + \alpha_2 f_0, \\ f_3' &= f_3 f_1 (f_0 - f_2) + \left(\frac{1}{2} - \alpha_1\right) f_3 + \alpha_3 f_1, \end{aligned} \qquad {}' = t\frac{d}{dt}, \tag{2.1}$$

with the normalization conditions

$$\alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 = 1, \quad f_0 + f_2 = f_1 + f_3 = \sqrt{t}. \tag{2.2}$$

The correspondence to the canonical variables and parameters of $\mathrm{S_V}$ is given by

$$q = -\frac{f_3}{f_1}, \quad p = \frac{1}{\sqrt{t}} f_1 (f_0 f_1 + \alpha_0), \tag{2.3}$$

and

$$\kappa_\infty = \alpha_1, \quad \kappa_0 = \alpha_3, \quad \theta = \alpha_2 - \alpha_0 - 1, \tag{2.4}$$

respectively. The Bäcklund transformations of $\mathrm{P_V}$ are described as follows,

$$\begin{aligned} & s_i(\alpha_i) = -\alpha_i, \quad s_i(\alpha_j) = \alpha_j + \alpha_i \ \ (j = i \pm 1), \quad s_i(\alpha_j) = \alpha_j \ \ (j \neq i, i \pm 1), \\ & s_i(f_i) = f_i, \qquad s_i(f_j) = f_j \pm \frac{\alpha_i}{f_i} \ \ (j = i \pm 1), \quad s_i(f_j) = f_j \ \ (j \neq i, i \pm 1), \\ & \pi(\alpha_j) = \alpha_{j+1}, \quad \pi(f_j) = f_{j+1}, \end{aligned} \tag{2.5}$$

where the subscripts $i = 0, 1, 2, 3$ are understood as elements of $\mathbb{Z}/4\mathbb{Z}$. The Hamiltonians h_i of the system (2.1) are given by

$$\begin{aligned} h_0 = f_0 f_1 f_2 f_3 &+ \frac{\alpha_1 + 2\alpha_2 - \alpha_3}{4} f_0 f_1 + \frac{\alpha_1 + 2\alpha_2 + 3\alpha_3}{4} f_1 f_2 \\ &- \frac{3\alpha_1 + 2\alpha_2 + \alpha_3}{4} f_2 f_3 + \frac{\alpha_1 - 2\alpha_2 - \alpha_3}{4} f_3 f_0 + \frac{(\alpha_1 + \alpha_3)^2}{4}, \end{aligned} \tag{2.6}$$

and $h_i = \pi^i(h_0)$. Then, we have

$$s_i(h_j) = h_j \ (i \neq j), \quad s_i(h_i) = h_i + \sqrt{t}\, \frac{\alpha_i}{f_i}, \quad \pi(h_i) = h_{i+1}. \tag{2.7}$$

Introducing the τ-functions τ_i as

$$h_i = (\log \tau_i)', \tag{2.8}$$

we find that the Bäcklund transformations for the τ-functions are described as

$$s_i(\tau_j) = \tau_j \quad (i \neq j), \quad s_i(\tau_i) = f_i \frac{\tau_{i-1}\tau_{i+1}}{\tau_i}, \quad \pi(\tau_i) = \tau_{i+1}. \tag{2.9}$$

The canonical variables of $\mathrm{S_V}$ are recovered from the τ-functions by

$$q = -\frac{\tau_3 s_3(\tau_3)}{\tau_1 s_1(\tau_1)}, \quad p = \frac{1}{\sqrt{t}} \frac{\tau_1 s_1(\tau_1) s_0 s_1(\tau_1)}{\tau_2^2 \tau_3}. \tag{2.10}$$

Let us define the translation operators T_i by

$$T_1 = \pi s_3 s_2 s_1, \quad T_2 = s_1 \pi s_3 s_2, \quad T_3 = s_2 s_1 \pi s_3, \quad T_0 = s_3 s_2 s_1 \pi, \tag{2.11}$$

which commute with each other and act on parameters α_i as

$$T_i(\alpha_{i-1}) = \alpha_{i-1} + 1, \quad T_i(\alpha_i) = \alpha_i - 1, \quad T_i(\alpha_j) = \alpha_j \ (j \neq i-1, i). \tag{2.12}$$

Noting that $T_1 T_2 T_3 T_0 = 1$, we set

$$\tau_{k,l,m} = T_1^k T_2^l T_3^m(\tau_0), \quad k, l, m \in \mathbb{Z}. \tag{2.13}$$

Then, from (2.9) and (2.10), we have

$$\begin{aligned} T_1^k T_2^l T_3^m(f_0) &= \frac{\tau_{k,l,m}\tau_{k+2,l+1,m+1}}{\tau_{k+1,l+1,m+1}\tau_{k+1,l,m}}, \quad & T_1^k T_2^l T_3^m(f_1) &= \frac{\tau_{k+1,l,m}\tau_{k,l+1,m}}{\tau_{k,l,m}\tau_{k+1,l+1,m}}, \\ T_1^k T_2^l T_3^m(f_2) &= \frac{\tau_{k+1,l+1,m}\tau_{k+1,l,m+1}}{\tau_{k+1,l,m}\tau_{k+1,l+1,m+1}}, \quad & T_1^k T_2^l T_3^m(f_3) &= \frac{\tau_{k+1,l+1,m+1}\tau_{k,l,m-1}}{\tau_{k+1,l+1,m}\tau_{k,l,m}}, \end{aligned} \tag{2.14}$$

and

$$T_1^k T_2^l T_3^m(q) = -\frac{\tau_{k+1,l+1,m+1}\tau_{k,l,m-1}}{\tau_{k+1,l,m}\tau_{k,l+1,m}}, \quad T_1^k T_2^l T_3^m(p) = \frac{1}{\sqrt{t}} \frac{\tau_{k+1,l,m}\tau_{k,l+1,m}\tau_{k+2,l+2,m+1}}{\tau_{k+1,l+1,m}^2 \tau_{k+1,l+1,m+1}}, \tag{2.15}$$

respectively. It is possible to derive the Toda equation with respect to each translation operator. For the T_1-direction, we have

$$\tau_{k+1,l,m}\tau_{k-1,l,m} = \frac{1}{\sqrt{t}} \left[(\log \tau_{k,l,m})'' + \frac{3\alpha_1 + 2\alpha_2 + \alpha_3 - 3k + l + m}{4} t \right] \tau_{k,l,m} \cdot \tau_{k,l,m}. \tag{2.16}$$

2.2 A Riccati solution

If $\alpha_0 = 0$, the right-hand side of the first equation of (2.1) is divisible by f_0. This means that the system (2.1) admits the specialization of $f_0 = 0$ if $\alpha_0 = 0$. Then, setting $f_1 = \sqrt{t} f$, we see that f satisfies a Riccati equation

$$f' = tf(1-f) - (\alpha_1 + \alpha_3)f + \alpha_1. \tag{2.17}$$

By the dependent variable transformation $f = \dfrac{d}{dt}\log\varphi$, we have for φ the linear equation

$$\left[t\frac{d^2}{dt^2} + (\alpha_1 + \alpha_3 - t)\frac{d}{dt} - \alpha_1\right]\varphi = 0, \tag{2.18}$$

which is nothing but the confluent hypergeometric differential equation. We set $\alpha_1 = a$ and $\alpha_3 = c - a$, and impose "non-integer conditions" on them appropriately. The general solution of (2.18) is expressed by

$$\varphi = c_1\frac{\Gamma(a)\Gamma(c-a)}{\Gamma(c)}F(a,c;t) + c_2\frac{1}{\sin\pi(c-a)\Gamma(2-c)}t^{1-c}F(a-c+1,2-c;t), \tag{2.19}$$

where $F(a,c;t)$ denotes Kummer's confluent hypergeometric function, and c_i $(i = 1,2)$ are arbitrary complex constants. For simplicity, we denote

$$f_{i,j} = F(a+i,c+j;t), \quad g_{i,j} = t^{1-c-j}F(a-c+1+i-j,2-c-j;t) \quad (i,j \in \mathbb{Z}). \tag{2.20}$$

By using the contiguity relations of Kummer's function, we obtain the following proposition.

Proposition 2.1 *We define the functions* $\varphi_{i,j}$ *by*

$$\varphi_{i,j} = c_1\frac{\Gamma(a+i)\Gamma(c-a-i+j)}{\Gamma(c+j)}f_{i,j} + c_2\frac{1}{\sin\pi(c-a-i+j)\Gamma(2-c-j)}g_{i,j}. \tag{2.21}$$

Then,

$$(f_0,f_1,f_2,f_3) = \left(0, \sqrt{t}\,\frac{\varphi_{1,1}}{\varphi_{0,0}}, \sqrt{t}, \sqrt{t}\frac{\varphi_{0,1}}{\varphi_{0,0}}\right), \quad (\alpha_0,\alpha_1,\alpha_2,\alpha_3) = (0,a,1-c,c-a), \tag{2.22}$$

and

$$q = -\frac{\varphi_{0,1}}{\varphi_{1,1}}, \quad p = 0, \qquad \kappa_\infty = a, \quad \kappa_0 = c-a, \quad \theta = -c, \tag{2.23}$$

give a Riccati solution of the symmetric form of $P_{\rm V}$ *and the Hamiltonian system* $S_{\rm V}$*, respectively.*

2.3 A Determinant formula for a family of classical transcendental solutions

First, we calculate the Hamiltonians and τ-functions for the Riccati solution in Proposition 2.1. Under the specialization (2.22), the Hamiltonians and τ-functions are calculated as

$$\begin{gathered} h_0 = \frac{\varphi_{0,0}'}{\varphi_{0,0}} - \frac{2a-c+2}{4}t + \frac{c^2}{4}, \quad h_1 = -\frac{2a-c-1}{4}t + \frac{(c-1)^2}{4}, \\ h_2 = -\frac{2a-c}{4}t + \frac{c^2}{4}, \quad h_3 = -\frac{2a-c+1}{4}t + \frac{(c-1)^2}{4}, \end{gathered} \tag{2.24}$$

and

$$\begin{aligned}
&\tau_0=\tau_{0,0,0}=\varphi_{0,0}\,t^{c^2/4}\exp\left(-\frac{2a-c+2}{4}t\right),\quad \tau_1=\tau_{1,0,0}=t^{(c-1)^2/4}\exp\left(-\frac{2a-c-1}{4}t\right),\\
&\tau_2=\tau_{1,1,0}=t^{c^2/4}\exp\left(-\frac{2a-c}{4}t\right),\quad \tau_3=\tau_{1,1,1}=t^{(c-1)^2/4}\exp\left(-\frac{2a-c+1}{4}t\right),\\
&s_0(\tau_0)=\tau_{2,1,1}=0,\quad s_1(\tau_1)=\tau_{0,1,0}=\varphi_{1,1}t^{(c+1)^2/4}\exp\left(-\frac{2a-c+3}{4}t\right),\\
&s_2(\tau_2)=\tau_{1,0,1}=t^{(c-2)^2/4}\exp\left(-\frac{2a-c}{4}t\right),\\
&s_3(\tau_3)=\tau_{0,0,-1}=\varphi_{0,1}t^{(c+1)^2/4}\exp\left(-\frac{2a-c+1}{4}t\right),
\end{aligned}\tag{2.25}$$

up to multiplication by some constants, respectively. For small k,l,m, we observe that $\tau_{k,l,m}$ are expressed in the form

$$\tau_{k,l,m}=\sigma_{k,l,m}t^{(c-k+l-m)^2/4-k(k-1)/2}\exp\left(-\frac{2a-c+2-3k+l+m}{4}t\right),\tag{2.26}$$

with

$$\sigma_{2,l,m}=0,\quad \sigma_{1,l,m}=\text{const.},\quad \sigma_{0,l,m}=(\text{const.})\times\varphi_{l,l-m}.\tag{2.27}$$

Assume that $\tau_{k,l,m}$ are expressed as (2.26) for any $k,l,m\in\mathbb{Z}$. Then, the Toda equation (2.16) yields

$$\sigma_{k+1,l,m}\sigma_{k-1,l,m}=\sigma''_{k,l,m}\sigma_{k,l,m}-(\sigma'_{k,l,m})^2.\tag{2.28}$$

Moreover, we set

$$\sigma_{k,l,m}=\omega_{k,l,m}\rho_{k,l,m},\quad \omega_{k,l,m}=\omega_{k,l,m}(a,c),\tag{2.29}$$

with $\rho_{1,l,m}=1$ and $\rho_{0,l,m}=\varphi_{l,l-m}$, and impose that the constants $\omega_{k,l,m}$ satisfy

$$\omega_{k+1,l,m}\omega_{k-1,l,m}=\omega_{k,l,m}^2.\tag{2.30}$$

Then, the functions $\rho_{k,l,m}$ are determined by the recurrence relation

$$\rho_{k+1,l,m}\rho_{k-1,l,m}=\rho''_{k,l,m}\rho_{k,l,m}-(\rho'_{k,l,m})^2,\tag{2.31}$$

with the initial conditions

$$\rho_{2,l,m}=0,\quad \rho_{1,l,m}=1,\quad \rho_{0,l,m}=\varphi_{l,l-m}.\tag{2.32}$$

By Darboux's formula, the functions $\rho_{1-n,l,m}$ for $n\in\mathbb{Z}_{\geq 0}$ are expressed as

$$\rho_{1-n,l,m}=\begin{vmatrix}\varphi_{l,l-m}^{(0)} & \varphi_{l,l-m}^{(1)} & \cdots & \varphi_{l,l-m}^{(n-1)}\\ \varphi_{l,l-m}^{(1)} & \varphi_{l,l-m}^{(2)} & \cdots & \varphi_{l,l-m}^{(n)}\\ \vdots & \vdots & \ddots & \vdots\\ \varphi_{l,l-m}^{(n-1)} & \varphi_{l,l-m}^{(n)} & \cdots & \varphi_{l,l-m}^{(2n-2)}\end{vmatrix},\quad \varphi_{l,l-m}^{(i)}=\left(t\frac{d}{dt}\right)^i\varphi_{l,l-m}.\tag{2.33}$$

Note that the constants $\omega_{k,l,m}$ are determined by the recurrence relations (2.30) and

$$\begin{aligned}
&\omega_{1,l+1,m}\omega_{1,l-1,m}=-(a+l-1)\omega_{1,l,m}^2,\quad \omega_{1,l,m+1}\omega_{1,l,m-1}=(c-a-m)\omega_{1,l,m}^2,\\
&\omega_{0,l+1,m}\omega_{0,l-1,m}=-(a+l-1)\omega_{0,l,m}^2,\quad \omega_{0,l,m+1}\omega_{0,l,m-1}=(c-a-1-m)\omega_{0,l,m}^2,
\end{aligned}\tag{2.34}$$

with the initial conditions

$$\begin{aligned} &\omega_{1,0,0}=\omega_{1,1,0}=\omega_{1,0,1}=\omega_{1,1,1}=1,\\ &\omega_{0,0,0}=\omega_{0,1,0}=\omega_{0,0,-1}=\omega_{0,1,-1}=1. \end{aligned} \tag{2.35}$$

Since it is possible to set $l = m = 0$ without loss of generality, we obtain the following theorem.

Theorem 2.2 *We define the functions $\tau_n^{i,j}$ by*

$$\tau_n^{i,j}=\begin{vmatrix} \varphi_{i,j}^{(0)} & \varphi_{i,j}^{(1)} & \cdots & \varphi_{i,j}^{(n-1)}\\ \varphi_{i,j}^{(1)} & \varphi_{i,j}^{(2)} & \cdots & \varphi_{i,j}^{(n)}\\ \vdots & \vdots & \ddots & \vdots\\ \varphi_{i,j}^{(n-1)} & \varphi_{i,j}^{(n)} & \cdots & \varphi_{i,j}^{(2n-2)} \end{vmatrix},\quad \varphi_{i,j}^{(k)}=\left(t\frac{d}{dt}\right)^k\varphi_{i,j}, \tag{2.36}$$

where $\varphi_{i,j}$ are given by (2.21). Then,

$$\begin{aligned} &f_0=\frac{1}{\sqrt{t}}\frac{\tau_{n+1}^{0,0}\tau_{n-1}^{1,0}}{\tau_n^{1,0}\tau_n^{0,0}},\quad f_1=\sqrt{t}\frac{\tau_n^{0,0}\tau_{n+1}^{1,1}}{\tau_{n+1}^{0,0}\tau_n^{1,1}},\\ &f_2=\sqrt{t}\frac{\tau_n^{1,1}\tau_n^{0,-1}}{\tau_n^{0,0}\tau_n^{1,0}},\quad f_3=\sqrt{t}\left(\frac{c-a-1}{c-a}\right)^n\frac{\tau_n^{1,0}\tau_{n+1}^{0,1}}{\tau_n^{1,1}\tau_{n+1}^{0,0}}, \end{aligned} \tag{2.37}$$

$$(\alpha_0,\alpha_1,\alpha_2,\alpha_3)=(-n,a+n,1-c,c-a), \tag{2.38}$$

and

$$q=-\left(\frac{c-a-1}{c-a}\right)^n\frac{\tau_n^{1,0}\tau_{n+1}^{0,1}}{\tau_n^{0,0}\tau_{n+1}^{1,1}},\quad p=-\frac{a}{c-a-1}\frac{1}{t}\frac{\tau_n^{0,0}\tau_{n+1}^{1,1}\tau_{n-1}^{2,1}}{\tau_n^{1,1}\tau_n^{1,1}\tau_n^{1,0}}, \tag{2.39}$$

$$\kappa_\infty=a+n,\quad \kappa_0=c-a,\quad \theta=-c+n, \tag{2.40}$$

give a family of classical transcendental solutions of the symmetric form of P_{V} and the Hamiltonian system S_{V}, respectively.

Remark 2.3 *Noting that we have from (2.7)*

$$\sqrt{t}\ \frac{\alpha_i}{f_i}=t\frac{d}{dt}\log\frac{s_i(\tau_i)}{\tau_i}, \tag{2.41}$$

which lead us to another expression of the solutions in Theorem 2.2. For example, we obtain

$$(a+n)\frac{\sqrt{t}}{f_1}=t\frac{d}{dt}\log\frac{\tau_{n+1}^{1,1}}{\tau_n^{0,0}}+c-t. \tag{2.42}$$

Remark 2.4 *The symmetric form of P_{V} admits the following symmetry σ:*

$$\begin{aligned} &\sigma(t)=-t,\\ &\sigma(f_0)=\sqrt{-1}f_2,\quad \sigma(f_2)=\sqrt{-1}f_0,\quad \sigma(f_1)=\sqrt{-1}f_1,\quad \sigma(f_3)=\sqrt{-1}f_3,\\ &\sigma(\alpha_0)=\alpha_2,\quad \sigma(\alpha_2)=\alpha_0,\quad \sigma(\alpha_1)=\alpha_1,\quad \sigma(\alpha_3)=\alpha_3. \end{aligned} \tag{2.43}$$

The application of σ to the family of solutions in Theorem 2.2 gives another family of solutions expressed in terms of $F(a,c;-t)$.

3 Classical transcendental solutions of the Painlevé VI equation

3.1 The symmetric form of the Painlevé VI equation

Here, we give a brief review of the symmetric form of the Painlevé VI equation [9, 4]. We set

$$f_0 = q - t, \quad f_3 = q - 1, \quad f_4 = q, \quad f_2 = p, \tag{3.1}$$

and

$$\alpha_0 = \theta, \quad \alpha_1 = \kappa_\infty, \quad \alpha_3 = \kappa_1, \quad \alpha_4 = \kappa_0. \tag{3.2}$$

Then, the Hamiltonian (1.6) is written as

$$H = f_2^2 f_0 f_3 f_4 - [(\alpha_0 - 1) f_3 f_4 + \alpha_3 f_0 f_4 + \alpha_4 f_0 f_3] f_2 + \alpha_2(\alpha_1 + \alpha_2) f_0, \tag{3.3}$$

with

$$\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = 1, \tag{3.4}$$

and the Hamilton equation (1.5) is written as

$$\begin{aligned} f_4' &= 2 f_2 f_0 f_3 f_4 - (\alpha_0 - 1) f_3 f_4 - \alpha_3 f_0 f_4 - \alpha_4 f_0 f_3, \\ f_2' &= -(f_0 f_3 + f_0 f_4 + f_3 f_4) f_2^2 \\ &\qquad + [(\alpha_0 - 1)(f_3 + f_4) + \alpha_3(f_0 + f_4) + \alpha_4(f_0 + f_3)] f_2 - \alpha_2(\alpha_1 + \alpha_2). \end{aligned} \tag{3.5}$$

The fundamental Bäcklund transformations of $\mathrm{P_{VI}}$ are given in Table 1. We define the Hamiltonians h_i $(i = 0, 1, 2, 3, 4)$ as

$$h_0 = H_0 + \frac{t}{4}, \quad h_1 = s_5(H_0) - \frac{t-1}{4}, \quad h_3 = s_6(H_0) + \frac{1}{4}, \quad h_4 = s_7(H_0), \quad h_2 = h_1 + s_1(h_1), \tag{3.6}$$

where an auxiliary Hamiltonian H_0 is given by

$$H_0 = H + \frac{t}{4}[1 + 4\alpha_1\alpha_2 + 4\alpha_2^2 - (\alpha_3 + \alpha_4)^2] + \frac{1}{4}[(\alpha_1 + \alpha_4)^2 + (\alpha_3 + \alpha_4)^2 + 4\alpha_2\alpha_4]. \tag{3.7}$$

Introducing the τ-functions τ_i as

$$h_i = (\log \tau_i)', \tag{3.8}$$

we find that the Bäcklund transformations for the τ-functions are described as follows,

$$\begin{aligned} &s_i(\tau_j) = \tau_j \quad (i \neq j, \;\; i, j = 0, 1, 2, 3, 4), \\ &s_i(\tau_i) = f_i \frac{\tau_2}{\tau_i} \quad (i = 0, 3, 4), \quad s_1(\tau_1) = \frac{\tau_2}{\tau_1}, \quad s_2(\tau_2) = \frac{f_2}{\sqrt{t}} \frac{\tau_0\tau_1\tau_3\tau_4}{\tau_2}, \end{aligned} \tag{3.9}$$

$$\begin{aligned} s_5: \quad &\tau_0 \to [t(t-1)]^{\frac{1}{4}}\tau_1, \qquad \tau_1 \to [t(t-1)]^{-\frac{1}{4}}\tau_0, \\ &\tau_3 \to t^{-\frac{1}{4}}(t-1)^{\frac{1}{4}}\tau_4, \qquad \tau_4 \to t^{\frac{1}{4}}(t-1)^{-\frac{1}{4}}\tau_3, \\ &\tau_2 \to [t(t-1)]^{-\frac{1}{2}} f_0 \tau_2, \end{aligned} \tag{3.10}$$

$$s_6: \quad \tau_0 \to i t^{\frac{1}{4}}\tau_3, \quad \tau_3 \to -i t^{-\frac{1}{4}}\tau_0, \quad \tau_1 \to t^{-\frac{1}{4}}\tau_4, \quad \tau_4 \to t^{\frac{1}{4}}\tau_1, \quad \tau_2 \to t^{-\frac{1}{2}} f_4 \tau_2, \tag{3.11}$$

$$\begin{aligned} s_7: \quad &\tau_0 \to (-1)^{-\frac{3}{4}}(t-1)^{\frac{1}{4}}\tau_4, \quad \tau_4 \to (-1)^{\frac{3}{4}}(t-1)^{-\frac{1}{4}}\tau_0, \\ &\tau_1 \to (-1)^{\frac{3}{4}}(t-1)^{-\frac{1}{4}}\tau_3, \quad \tau_3 \to (-1)^{-\frac{3}{4}}(t-1)^{\frac{1}{4}}\tau_1, \\ &\tau_2 \to -i(t-1)^{-\frac{1}{2}} f_3 \tau_2. \end{aligned} \tag{3.12}$$

Table 1: Bäcklund transformations of $\mathrm{P_{VI}}$

	α_0	α_1	α_2	α_3	α_4	f_4	f_2
s_0	$-\alpha_0$	α_1	$\alpha_2+\alpha_0$	α_3	α_4	f_4	$f_2-\dfrac{\alpha_0}{f_0}$
s_1	α_0	$-\alpha_1$	$\alpha_2+\alpha_1$	α_3	α_4	f_4	f_2
s_2	$\alpha_0+\alpha_2$	$\alpha_1+\alpha_2$	$-\alpha_2$	$\alpha_3+\alpha_2$	$\alpha_4+\alpha_2$	$f_4+\dfrac{\alpha_2}{f_2}$	f_2
s_3	α_0	α_1	$\alpha_2+\alpha_3$	$-\alpha_3$	α_4	f_4	$f_2-\dfrac{\alpha_3}{f_3}$
s_4	α_0	α_1	$\alpha_2+\alpha_4$	α_3	$-\alpha_4$	f_4	$f_2-\dfrac{\alpha_4}{f_4}$
s_5	α_1	α_0	α_2	α_4	α_3	$t\dfrac{f_3}{f_0}$	$-\dfrac{f_0(f_2f_0+\alpha_2)}{t(t-1)}$
s_6	α_3	α_4	α_2	α_0	α_1	$\dfrac{t}{f_4}$	$-\dfrac{f_4(f_4f_2+\alpha_2)}{t}$
s_7	α_4	α_3	α_2	α_1	α_0	$\dfrac{f_0}{f_3}$	$\dfrac{f_3(f_3f_2+\alpha_2)}{t-1}$

Let us define the following translation operators

$$\begin{aligned} \widehat{T}_{13} &= s_1s_2s_0s_4s_2s_1s_7, \quad \widehat{T}_{40} = s_4s_2s_1s_3s_2s_4s_7, \\ \widehat{T}_{34} &= s_3s_2s_0s_1s_2s_3s_5, \quad T_{14} = s_1s_4s_2s_0s_3s_2s_6, \end{aligned} \tag{3.13}$$

which act on parameters α_i as

$$\begin{aligned} \widehat{T}_{13}(\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4) &= (\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4)+(0,1,0,-1,0), \\ \widehat{T}_{40}(\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4) &= (\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4)+(-1,0,0,0,1), \\ \widehat{T}_{34}(\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4) &= (\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4)+(0,0,0,1,-1), \\ T_{14}(\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4) &= (\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4)+(0,1,-1,0,1). \end{aligned} \tag{3.14}$$

Note that the action of these operators on τ-functions is not commutative. For example, we have

$$\begin{aligned} &\widehat{T}_{13}\widehat{T}_{40}(\tau_0) = -\widehat{T}_{40}\widehat{T}_{13}(\tau_0), \quad \widehat{T}_{13}\widehat{T}_{34}(\tau_0) = -i\widehat{T}_{34}\widehat{T}_{13}(\tau_0), \\ &\widehat{T}_{13}T_{14}(\tau_0) = iT_{14}\widehat{T}_{13}(\tau_0), \quad \widehat{T}_{40}\widehat{T}_{34}(\tau_0) = i\widehat{T}_{34}\widehat{T}_{40}(\tau_0), \\ &\widehat{T}_{40}T_{14}(\tau_0) = -iT_{14}\widehat{T}_{40}(\tau_0), \quad \widehat{T}_{34}T_{14}(\tau_0) = iT_{14}\widehat{T}_{34}(\tau_0). \end{aligned} \tag{3.15}$$

Setting

$$\tau_{k,l,m,n} = T_{14}^n\widehat{T}_{34}^m\widehat{T}_{40}^l\widehat{T}_{13}^k(\tau_0), \quad k,l,m,n\in\mathbb{Z}, \tag{3.16}$$

we have

$$\begin{aligned} T_{14}^n\widehat{T}_{34}^m\widehat{T}_{40}^l\widehat{T}_{13}^k(f_0) &= -it^{\frac{1}{2}}(t-1)^{\frac{1}{2}}\frac{\tau_{k,l,m,n}\tau_{k-1,l-2,m-1,n+1}}{\tau_{k-1,l-1,m-1,n}\tau_{k,l-1,m,n+1}}, \\ T_{14}^n\widehat{T}_{34}^m\widehat{T}_{40}^l\widehat{T}_{13}^k(f_3) &= i(t-1)^{\frac{1}{2}}\frac{\tau_{k,l-1,m-1,n}\tau_{k-1,l-1,m,n+1}}{\tau_{k-1,l-1,m-1,n}\tau_{k,l-1,m,n+1}}, \\ T_{14}^n\widehat{T}_{34}^m\widehat{T}_{40}^l\widehat{T}_{13}^k(f_4) &= t^{\frac{1}{2}}\frac{\tau_{k,l-1,m,n}\tau_{k-1,l-1,m-1,n+1}}{\tau_{k-1,l-1,m-1,n}\tau_{k,l-1,m,n+1}}, \\ T_{14}^n\widehat{T}_{34}^m\widehat{T}_{40}^l\widehat{T}_{13}^k(f_2) &= -(t-1)^{-\frac{1}{2}}\frac{\tau_{k-1,l-1,m-1,n}\tau_{k,l-1,m,n+1}\tau_{k+1,l,m,n-1}}{\tau_{k,l,m,n}\tau_{k,l-1,m-1,n}\tau_{k,l-1,m,n}}. \end{aligned} \tag{3.17}$$

It is possible to derive the Toda equation with respect to each translation operator. For the T_{14}-direction, we get

$$\begin{aligned}&\tau_{k,l,m,n+1}\tau_{k,l,m,n-1}\\&\quad=-t^{-\frac{1}{2}}\left[(t-1)\frac{d}{dt}(\log\tau_{k,l,m,n})'-(\log\tau_{k,l,m,n})'+\frac{(\alpha_1+\alpha_4+k+l-m+2n)^2}{4}+\frac{1}{2}\right]\tau_{k,l,m,n}^2.\end{aligned}\tag{3.18}$$

3.2 A Riccati solution

If $\alpha_2=0$, it is possible to specialize $f_2=0$. Then, the Hamilton equation (3.5) yields

$$f_4'=-(\alpha_0-1)f_3f_4-\alpha_3f_0f_4-\alpha_4f_0f_3,\tag{3.19}$$

which is equivalent to a Riccati equation

$$q'=\alpha_1q^2+[(\alpha_3+\alpha_4)t+(\alpha_0+\alpha_4-1)]q-\alpha_4t.\tag{3.20}$$

We set

$$\alpha_0=-b,\quad \alpha_1=a,\quad \alpha_3=c-a,\quad \alpha_4=b-c+1,\tag{3.21}$$

and impose "non-integer conditions" on them appropriately. By the dependent variable transformation

$$aq=-\frac{\varphi'}{\varphi}-(b+1)t+c,\tag{3.22}$$

we have for φ the linear equation

$$t(t-1)\frac{d^2\varphi}{dt^2}+[(a+b+3)t-(c+1)]\frac{d\varphi}{dt}+(a+1)(b+1)\varphi=0,\tag{3.23}$$

which is nothing but the hypergeometric differential equation. The general solution of (3.23) is expressed by

$$\begin{aligned}\varphi=\;&c_1\frac{\Gamma(a+1)\Gamma(b+1)}{\Gamma(c+1)}F(a+1,b+1,c+1;t)\\&+c_2\frac{\Gamma(a-c+1)\Gamma(b-c+1)}{\Gamma(1-c)}t^{-c}F(a-c+1,b-c+1,1-c;t),\end{aligned}\tag{3.24}$$

where $F(a,b,c;t)$ denotes Gauss's hypergeometric function and c_i $(i=1,2)$ are arbitrary complex constants. For simplicity, we denote

$$\left.\begin{aligned}f_m^{kl}&=F(a+k,b+l,c+m;t),\\g_m^{kl}&=t^{1-c-m}F(a-c+1+k-m,b-c+1+l-m,2-c-m;t)\end{aligned}\right.\qquad(k,l,m\in\mathbb{Z}).\tag{3.25}$$

By using the contiguity relations of Gauss's hypergeometric function, we obtain the following proposition.

Proposition 3.1 *We define the function* $\varphi_{k,l,m}$ *by*

$$\begin{aligned}\varphi_{k,l,m}=\;&c_1\frac{\Gamma(a+k+1)\Gamma(b+l+2)}{\Gamma(c+m+1)}f_{m+1}^{k+1,l+2}\\&+c_2\frac{\Gamma(a-c+1+k-m)\Gamma(b-c+2+l-m)}{\Gamma(1-c-m)}g_{m+1}^{k+1,l+2}.\end{aligned}\tag{3.26}$$

Then,

$$\begin{gathered}f_0=b\frac{\varphi_{-1,-2,-1}}{\varphi_{0,-1,0}},\quad f_3=(c-a)\frac{\varphi_{-1,-1,0}}{\varphi_{0,-1,0}},\quad f_4=\frac{\varphi_{-1,-1,-1}}{\varphi_{0,-1,0}},\quad f_2=0,\\\alpha_0=-b,\quad\alpha_1=a,\quad\alpha_3=c-a,\quad\alpha_4=b-c+1,\end{gathered}\tag{3.27}$$

gives a Riccati solution of the symmetric form of $P_{\rm VI}$.

3.3 A Determinant formula for a family of classical transcendental solutions

First, we calculate the Hamiltonians and τ-functions for the Riccati solution in Proposition 3.1. Under the specialization (3.27), the Hamiltonians are calculated as

$$h_i = A_i(t-1) + B_i t, \quad (i = 0, 1, 3, 4), \tag{3.28}$$

with

$$\begin{aligned} A_0 &= -\tfrac{1}{4}(a+b-c+1)^2 - \tfrac{1}{4}(a-b-1)^2, & B_0 &= \tfrac{1}{4}(a+b-c+1)^2 + \tfrac{1}{2}, \\ A_1 &= -\tfrac{1}{4}(a+b-c)^2 - \tfrac{1}{4}(a-b-1)^2 - \tfrac{1}{4}, & B_1 &= \tfrac{1}{4}(a+b-c)^2 + \tfrac{1}{4}, \\ A_3 &= -\tfrac{1}{4}(a+b-c+1)^2 - \tfrac{1}{4}(a-b)^2 - \tfrac{1}{4}, & B_3 &= \tfrac{1}{4}(a+b-c+1)^2 + \tfrac{1}{2}, \\ A_4 &= -\tfrac{1}{4}(a+b-c)^2 - \tfrac{1}{4}(a-b)^2, & B_4 &= \tfrac{1}{4}(a+b-c)^2 + \tfrac{1}{4}. \end{aligned} \tag{3.29}$$

Then, we have

$$\tau_i = t^{A_i}(t-1)^{B_i}, \quad s_2 s_i(\tau_i) = 0 \quad (i = 0, 1, 3, 4), \tag{3.30}$$

$$\begin{aligned} s_0(\tau_0) &= b\varphi_{-1,-2,-1} t^{A_0+b}(t-1)^{B_0}, & s_1(\tau_1) &= \varphi_{0,-1,0} t^{A_1+(c-a)}(t-1)^{B_1+(a+b-c+1)}, \\ s_3(\tau_3) &= (c-a)\varphi_{-1,-1,0} t^{A_3+a}(t-1)^{B_3}, & s_4(\tau_4) &= \varphi_{-1,-1,-1} t^{A_4-(b-c+1)}(t-1)^{B_4+(a+b-c+1)}, \end{aligned} \tag{3.31}$$

up to multiplication by some constants. For small k, l, m, n, we observe that $\tau_{k,l,m,n}$ are expressed in the form

$$\tau_{k,l,m,n} = \sigma_{k,l,m,n} t^{-\frac{1}{4}(\hat{a}+\hat{b}-\hat{c}+2n)^2 - \frac{1}{4}(\hat{a}-\hat{b}-n)^2 + n(\hat{b}+n) - \frac{1}{2}n(n-1)} (t-1)^{\frac{1}{4}(\hat{a}+\hat{b}-\hat{c}+2n)^2 + \frac{1}{2}}, \tag{3.32}$$

with

$$\sigma_{k,l,m,-1} = 0, \quad \sigma_{k,l,m,0} = \text{const.}, \quad \sigma_{k,l,m,1} = (\text{const.}) \times \varphi_{k,l,m}, \tag{3.33}$$

where we denote

$$\hat{a} = a + k, \quad \hat{b} = b + l + 1, \quad \hat{c} = c + m. \tag{3.34}$$

Assume that $\tau_{k,l,m,n}$ are expressed as (3.32) for any $k, l, m, n \in \mathbb{Z}$. Then, the Toda equation (3.18) yields

$$\sigma_{k,l,m,n+1}\sigma_{k,l,m,n-1} = -\left[(\delta^2\sigma_{k,l,m,n})\sigma_{k,l,m,n} - (\delta\sigma_{k,l,m,n})^2\right], \quad \delta = t\frac{d}{dt}. \tag{3.35}$$

Moreover, we set

$$\sigma_{k,l,m,n} = \omega_{k,l,m,n}\rho_{k,l,m,n}, \quad \omega_{k,l,m,n} = \omega_{k,l,m,n}(a, b, c), \tag{3.36}$$

with $\rho_{k,l,m,0} = 1$ and $\rho_{k,l,m,1} = \varphi_{k,l,m}$, and impose that the constants $\omega_{k,l,m,n}$ satisfy

$$\omega_{k,l,m,n+1}\omega_{k,l,m,n-1} = -\omega_{k,l,m,n}^2. \tag{3.37}$$

Then, the function $\rho_{k,l,m,n}$ are determined by the recurrence relation

$$\rho_{k,l,m,n+1}\rho_{k,l,m,n-1} = (\delta^2\rho_{k,l,m,n})\rho_{k,l,m,n} - (\delta\rho_{k,l,m,n})^2, \tag{3.38}$$

with the initial conditions

$$\rho_{k,l,m,-1} = 0, \quad \rho_{k,l,m,0} = 1, \quad \rho_{k,l,m,1} = \varphi_{k,l,m}. \tag{3.39}$$

By Darboux's formula, the functions $\rho_{k,l,m,n}$ for $n \in \mathbb{Z}_{\geq 0}$ are expressed as

$$\rho_{k,l,m,n} = \begin{vmatrix} \varphi_{k,l,m}^{(0)} & \varphi_{k,l,m}^{(1)} & \cdots & \varphi_{k,l,m}^{(n-1)} \\ \varphi_{k,l,m}^{(1)} & \varphi_{k,l,m}^{(2)} & \cdots & \varphi_{k,l,m}^{(n)} \\ \vdots & \vdots & \ddots & \vdots \\ \varphi_{k,l,m}^{(n-1)} & \varphi_{k,l,m}^{(n)} & \cdots & \varphi_{k,l,m}^{(2n-2)} \end{vmatrix}, \quad \varphi_{k,l,m}^{(i)} = \left(t\frac{d}{dt}\right)^i \varphi_{k,l,m}. \tag{3.40}$$

Note that the constants $\omega_{k,l,m,n}$ are determined by the recurrence relations (3.37) and

$$\begin{aligned}
\omega_{k+1,l,m,i}\omega_{k-1,l,m,i} &= i\hat{a}(\hat{c}-\hat{a})\omega_{k,l,m,i}^2, \\
\omega_{k,l+1,m,i}\omega_{k,l-1,m,i} &= -i\hat{b}(\hat{c}-\hat{b})\omega_{k,l,m,i}^2, \qquad (i=0,1) \\
\omega_{k,l,m+1,i}\omega_{k,l,m-1,i} &= (\hat{c}-\hat{a})(\hat{c}-\hat{b})\omega_{k,l,m,i}^2,
\end{aligned} \tag{3.41}$$

with the initial conditions

$$\begin{gathered}
\omega_{-1,-2,-1,1} = (-1)^{-\frac{1}{4}}b, \quad \omega_{0,-2,-1,1} = b, \quad \omega_{-1,-1,-1,1} = 1, \quad \omega_{0,-1,-1,1} = (-1)^{-\frac{1}{4}}, \\
\omega_{-1,0,0,1} = -(-1)^{-\frac{3}{4}}(c-a), \quad \omega_{0,0,0,1} = -i, \quad \omega_{-1,-1,0,1} = -i(c-a), \quad \omega_{0,-1,0,1} = (-1)^{-\frac{3}{4}},
\end{gathered} \tag{3.42}$$

and

$$\begin{gathered}
\omega_{-1,-2,-1,0} = (-1)^{-\frac{3}{4}}b, \quad \omega_{0,-2,-1,0} = -b, \quad \omega_{-1,-1,-1,0} = 1, \quad \omega_{0,-1,-1,0} = (-1)^{-\frac{3}{4}}, \\
\omega_{-1,0,0,0} = (-1)^{-\frac{3}{4}}(c-a), \quad \omega_{0,0,0,0} = 1, \quad \omega_{-1,-1,0,0} = c-a, \quad \omega_{0,-1,0,0} = (-1)^{-\frac{3}{4}}.
\end{gathered} \tag{3.43}$$

Since it is possible to set $k = l = m = 0$ without loss of generality, we obtain the following theorem.

Theorem 3.2 *We define the functions* $\tau_n^{k,l,m}$ *by*

$$\tau_n^{k,l,m} = \begin{vmatrix} \varphi_{k,l,m}^{(0)} & \varphi_{k,l,m}^{(1)} & \cdots & \varphi_{k,l,m}^{(n-1)} \\ \varphi_{k,l,m}^{(1)} & \varphi_{k,l,m}^{(2)} & \cdots & \varphi_{k,l,m}^{(n)} \\ \vdots & \vdots & \ddots & \vdots \\ \varphi_{k,l,m}^{(n-1)} & \varphi_{k,l,m}^{(n)} & \cdots & \varphi_{k,l,m}^{(2n-2)} \end{vmatrix}, \quad \varphi_{k,l,m}^{(i)} = \left(t\frac{d}{dt}\right)^i \varphi_{k,l,m}, \tag{3.44}$$

where $\varphi_{k,l,m}$ *are given by (3.26). Then,*

$$\begin{aligned}
f_0 &= b\frac{\tau_n^{0,0,0}\tau_{n+1}^{-1,-2,-1}}{\tau_n^{-1,-1,-1}\tau_{n+1}^{0,-1,0}}, \quad f_3 = (c-a)\frac{\tau_n^{0,-1,-1}\tau_{n+1}^{-1,-1,0}}{\tau_n^{-1,-1,-1}\tau_{n+1}^{0,-1,0}}, \\
f_4 &= \frac{\tau_n^{0,-1,0}\tau_{n+1}^{-1,-1,-1}}{\tau_n^{-1,-1,-1}\tau_{n+1}^{0,-1,0}}, \quad f_2 = at^{-1}\frac{\tau_n^{-1,-1,-1}\tau_{n+1}^{0,-1,0}\tau_{n-1}^{1,0,0}}{\tau_n^{0,0,0}\tau_n^{0,-1,-1}\tau_n^{0,-1,0}},
\end{aligned} \tag{3.45}$$

$$(\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4) = (-b, a+n, -n, c-a, b-c+1+n), \tag{3.46}$$

gives a family of classical transcendental solutions of the symmetric form of P_{VI}.

4 Degeneration of classical transcendental solutions

It is well known that, starting from $\mathrm{P_{VI}}$, one can obtain $\mathrm{P_V}, \ldots, \mathrm{P_I}$ by successive limiting procedures in the following diagram [14, 2],

$$\begin{tikzcd}
\mathrm{P_{VI}} \arrow[r] & \mathrm{P_V} \arrow[r] \arrow[d] & \mathrm{P_{III}} \arrow[d] & \\
& \mathrm{P_{IV}} \arrow[r] & \mathrm{P_{II}} \arrow[r] & \mathrm{P_I},
\end{tikzcd} \tag{4.1}$$

which corresponds to the degeneration diagram of the special functions of hypergeometric type

$$\begin{tikzcd}
\text{Gauss} \arrow[r] & \text{Kummer} \arrow[r] \arrow[d] & \text{Bessel} \arrow[d] \\
& \text{Hermite-Weber} \arrow[r] & \text{Airy}.
\end{tikzcd} \tag{4.2}$$

In this section, we show that, starting from the family of special function solutions of $\mathrm{P_{VI}}$ given in Theorem 3.2, we obtain classical transcendental solutions to other Painlevé equations by degeneration.

4.1 From $\mathrm{P_{VI}}$ to $\mathrm{P_V}$

As is known, the Hamiltonian system $\mathrm{S_{VI}}$ is reduced to $\mathrm{S_V}$ by putting

$$t \to 1-\varepsilon t, \quad \kappa_1 \to \varepsilon^{-1}+\theta+1, \quad \theta \to -\varepsilon^{-1}, \tag{4.3}$$

and taking the limit as $\varepsilon \to 0$. Let us consider the degeneration of the family of classical transcendental solutions given in Theorem 3.2. It is known that $\mathrm{P_{VI}}$ admits an outer symmetry as

$$\sigma_{34}: \quad \alpha_3 \leftrightarrow \alpha_4, \quad t \to 1-t, \quad f_4 \to -f_3, \quad f_2 \to -f_2. \tag{4.4}$$

The application of σ_{34} to the family of solutions in Theorem 3.2 gives another family of solutions for the parameters

$$(\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4) = (-b, a+n, -n, b-c+1+n, c-a), \tag{4.5}$$

or

$$\kappa_\infty = a+n, \quad \kappa_0 = c-a, \quad \kappa_1 = b-c+1+n, \quad \theta = -b. \tag{4.6}$$

Then, it is easy to see that by putting

$$t \to \varepsilon t, \quad b = \varepsilon^{-1}, \tag{4.7}$$

$\mathrm{S_{VI}}$ with (4.6) is reduced to $\mathrm{S_V}$ with

$$\kappa_\infty = a+n, \quad \kappa_0 = c-a, \quad \theta = -c+n, \tag{4.8}$$

in the limit as $\varepsilon \to 0$. It is obvious that Gauss's function $F(a,b,c;t)$ is reduced to Kummer's function $F(a,c;t)$ by this process. Thus, we get

$$\varphi_{k,l,m} \to \varepsilon^{-l}\Gamma^{-1}(c-a-k+m)\varphi_{k+1,m+1}, \tag{4.9}$$

where $\varphi_{i,j}$ are given by (2.21). Note that we redefine the constants c_1 and c_2 appropriately. It is easy to see that we have

$$\tau_n^{k,l,m} \to \varepsilon^{-ln}\Gamma^{-n}(c-a-k+m)\tau_n^{k+1,m+1}. \tag{4.10}$$

Therefore, we obtain the family of classical transcendental solutions of $\mathrm{S_V}$ in Theorem 2.2.

4.2 From $\mathrm{P_V}$ to $\mathrm{P_{III}}$

From $\mathrm{P_V}$, we can obtain two coalescence limits. First, we consider the Painlevé III equation

$$\frac{d^2q}{dt^2} = \frac{1}{q}\left(\frac{dq}{dt}\right)^2 - \frac{1}{t}\frac{dq}{dt} - \frac{4}{t}\left[\eta_\infty\theta_\infty q^2 + \eta_0(\theta_0+1)\right] + 4\eta_\infty^2 q^3 - \frac{4\eta_0^2}{q}, \tag{4.11}$$

which is equivalent to the Hamiltonian system

$$\mathrm{S_{III}}: \qquad q' = \frac{\partial H}{\partial p}, \quad p' = -\frac{\partial H}{\partial q}, \quad ' = t\frac{d}{dt}, \tag{4.12}$$

with the Hamiltonian

$$H = 2q^2p^2 - \left[2\eta_\infty t q^2 + (2\theta_0+1)q + 2\eta_0 t\right]p + \eta_\infty(\theta_0+\theta_\infty)tq. \tag{4.13}$$

This system can be derived from $\mathrm{S_V}$ by putting

$$\begin{gathered} q \to 1+\varepsilon t q, \quad p \to \varepsilon^{-1}t^{-1}p, \quad t \to \eta_0\varepsilon t^2, \quad H \to \frac{1}{2}(H+qp) \\ \kappa_\infty \to -\eta_\infty\varepsilon^{-1}+\theta_\infty, \quad \kappa_0 \to \eta_\infty\varepsilon^{-1}, \quad \theta \to \theta_0, \end{gathered} \tag{4.14}$$

and taking the limit as $\varepsilon \to 0$.

Let us apply the limiting procedure to the family of classical transcendental solutions of $\mathrm{S_V}$ given in Theorem 2.2. It is easy to see that by putting

$$a \to -\eta_\infty \varepsilon^{-1} + c, \tag{4.15}$$

$\mathrm{S_V}$ with (2.40) is reduced to $\mathrm{S_{III}}$ with

$$\theta_\infty = \nu + n + 1, \quad \theta_0 = -\nu + n - 1, \tag{4.16}$$

in the limit as $\varepsilon \to 0$, where we denote $c = \nu + 1$. Without loss of generality, it is possible to set

$$\eta_\infty = \frac{\sigma}{2}, \quad \eta_0 = \frac{1}{2}, \quad (\sigma = \pm 1). \tag{4.17}$$

Then, we find that

$$F(a, c; t) \to \Gamma(\nu + 1) \left(\frac{t}{2}\right)^{-\nu} Z_\nu(t), \tag{4.18}$$

where $Z_\nu = Z_\nu(t)$ denotes

$$Z_\nu = \begin{cases} J_\nu : & \text{Bessel} & \sigma = +1, \\ I_\nu : & \text{modified Bessel} & \sigma = -1. \end{cases} \tag{4.19}$$

This means that Kummer's function is reduced to the (modified) Bessel function in this limit. Thus, we get

$$\varphi_{i,j} \to (-1)^i \left(\frac{\sigma}{2}\varepsilon^{-1}\right)^j \left(\frac{t}{2}\right)^{-\nu-j} \varphi_{\nu+j}, \tag{4.20}$$

with

$$\varphi_{\nu+j} = c_1 Z_{\nu+j} + c_2 (-\sigma)^j Z_{-\nu-j}. \tag{4.21}$$

This leads us to

$$\tau_n^{i,j} \to \left(\frac{1}{2}\right)^{(n-1)n} (-1)^{in} \left(\frac{\sigma}{2}\varepsilon^{-1}\right)^{jn} \left(\frac{t}{2}\right)^{-(\nu+j)n} \tau_n^{\nu+j}, \tag{4.22}$$

where $\tau_n^{\nu+j}$ are defined by

$$\tau_n^{\nu+j} = \begin{vmatrix} \varphi_{\nu+j}^{(0)} & \varphi_{\nu+j}^{(1)} & \cdots & \varphi_{\nu+j}^{(n-1)} \\ \varphi_{\nu+j}^{(1)} & \varphi_{\nu+j}^{(2)} & \cdots & \varphi_{\nu+j}^{(n)} \\ \vdots & \vdots & \ddots & \vdots \\ \varphi_{\nu+j}^{(n-1)} & \varphi_{\nu+j}^{(n)} & \cdots & \varphi_{\nu+j}^{(2n-2)} \end{vmatrix}, \quad \varphi_{\nu+j}^{(i)} = \left(t\frac{d}{dt}\right)^i \varphi_{\nu+j}. \tag{4.23}$$

Therefore, from Theorem 2.2 and Remark 2.3, we obtain the following proposition [13].

Proposition 4.1 *(Okamoto) We define the functions τ_n^ν by (4.23). Then,*

$$q = \sigma \frac{\tau_{n+1}^\nu \tau_n^{\nu+1}}{\tau_n^\nu \tau_{n+1}^{\nu+1}} = \sigma \left(\frac{d}{dt} \log \frac{\tau_{n+1}^{\nu+1}}{\tau_n^\nu} + \frac{\nu + 1 - n}{t}\right), \quad p = -\frac{1}{2t} \frac{\tau_{n+1}^{\nu+1} \tau_{n-1}^{\nu+1}}{\tau_n^{\nu+1} \tau_n^{\nu+1}}, \tag{4.24}$$

$$\theta_\infty = \nu + n + 1, \quad \theta_0 = -\nu + n - 1, \tag{4.25}$$

with (4.17) gives a family of classical transcendental solutions of S_{III}.

4.3 From $\mathrm{P_V}$ to $\mathrm{P_{IV}}$

Next, we consider the Painlevé IV equation

$$\frac{d^2q}{dt^2} = \frac{1}{2q}\left(\frac{dq}{dt}\right)^2 + \frac{3}{2}q^3 + 2tq^2 + \frac{1}{2}t^2q - (-\kappa_0 + 2\theta_\infty + 1)q - \frac{\kappa_0^2}{2q}, \tag{4.26}$$

which is equivalent to the Hamiltonian system

$$\mathrm{S_{IV}}: \qquad q' = \frac{\partial H}{\partial p}, \quad p' = -\frac{\partial H}{\partial q}, \quad ' = \frac{d}{dt}, \tag{4.27}$$

with the Hamiltonian

$$H = qp^2 - (q^2 + tq + \kappa_0)p + \theta_\infty q. \tag{4.28}$$

This system can be also derived from $\mathrm{S_V}$ by coalescence. This process is achieved by putting

$$\begin{gathered} q \to \varepsilon q, \quad p \to \varepsilon^{-1}p, \quad t \to \varepsilon^{-2}(1 + \varepsilon t), \quad H + \kappa \to \varepsilon^{-1}H, \\ \theta \to \varepsilon^{-2} + 2\theta_\infty - \kappa_0, \quad \kappa_\infty \to \varepsilon^{-2}, \end{gathered} \tag{4.29}$$

and taking the limit as $\varepsilon \to 0$.

Let us consider the degeneration of the classical transcendental solutions of $\mathrm{S_V}$. Applying the Bäcklund transformation π^2 to the solutions in Theorem 2.2, we obtain the following corollary.

Corollary 4.2 *We define the functions* $\tau_n^{i,j}$ *by (2.36). Then,*

$$q = -\left(\frac{c-a}{c-a-1}\right)^n \frac{\tau_n^{0,0}\tau_{n+1}^{1,1}}{\tau_n^{1,0}\tau_{n+1}^{0,1}}, \quad p = (1-a)\left(\frac{c-a-1}{c-a}\right)^n \frac{\tau_n^{1,0}\tau_{n+1}^{0,1}\tau_{n+1}^{-1,-1}}{\tau_{n+1}^{0,0}\tau_{n+1}^{0,0}\tau_n^{0,0}}, \tag{4.30}$$

$$\kappa_\infty = c - a, \quad \kappa_0 = a + n, \quad \theta = c - n - 2, \tag{4.31}$$

gives a family of classical transcendental solutions of S_{V}.

It is easy to see that by putting

$$c \to \varepsilon^{-2} + a, \tag{4.32}$$

$\mathrm{S_V}$ with (4.31) is reduced to $\mathrm{S_{IV}}$ with

$$\kappa_0 = -\nu + n, \quad \theta_\infty = -\nu - 1, \tag{4.33}$$

in the limit as $\varepsilon \to 0$, where we denote $a = -\nu$. We find that

$$\frac{\Gamma(c-a)}{\Gamma(c)}F(a, c; t) \to (-\varepsilon)^{-\nu}H_\nu(t), \tag{4.34}$$

where $H_\nu(t)$ denotes the Hermite-Weber function. By a Kummer transformation

$$F(a - c + 1, 2 - c; t) = e^t F(1 - a, 2 - c; -t), \tag{4.35}$$

we get

$$\frac{\Gamma(c-a)\Gamma(a-c+1)}{\Gamma(2-c)}t^{1-c}F(a - c + 1, 2 - c; t) \to (-i\varepsilon)^{-\nu-1}e^{\frac{t^2}{2}}H_{-\nu-1}(it). \tag{4.36}$$

Thus, we have

$$\varphi_{k,j} \to (-\varepsilon)^{-\nu+k}\left[c_1\Gamma(-\nu + k)H_{\nu-k}(t) + c_2e^{i\pi(-\nu+k-1)/2}e^{\frac{t^2}{2}}H_{-\nu+k-1}(it)\right]. \tag{4.37}$$

Let us rewrite this expression in terms of the hyperbolic cylinder function $D_\nu(t)$. Noting the relations

$$H_\nu(t) = e^{\frac{t^2}{4}} D_\nu(t), \tag{4.38}$$

and

$$D_{-\nu-1}(it) = \frac{\Gamma(-\nu)}{\sqrt{2\pi}} \left[e^{i\pi(\nu+1)/2} D_\nu(t) - e^{-i\pi(\nu-1)/2} D_\nu(-t) \right], \tag{4.39}$$

we get

$$\varphi_{k,j} \to \varepsilon^k e^{\frac{t^2}{4}} \varphi_{\nu-k}, \tag{4.40}$$

with

$$\varphi_{\nu-k} = c_1 \frac{D_{\nu-k}(t)}{\Gamma(\nu-k+1)} + c_2 \Gamma(-\nu+k) D_{\nu-k}(-t). \tag{4.41}$$

This leads us to

$$\tau_n^{k,j} \to \varepsilon^{-n(n-1)} \varepsilon^{kn} \tau_n^{\nu-k}, \tag{4.42}$$

where $\tau_n^{\nu-k}$ are defined by

$$\tau_n^{\nu-k} = \begin{vmatrix} \phi_{\nu-k}^{(0)} & \phi_{\nu-k}^{(1)} & \cdots & \phi_{\nu-k}^{(n-1)} \\ \phi_{\nu-k}^{(1)} & \phi_{\nu-k}^{(2)} & \cdots & \phi_{\nu-k}^{(n)} \\ \vdots & \vdots & \ddots & \vdots \\ \phi_{\nu-k}^{(n-1)} & \phi_{\nu-k}^{(n)} & \cdots & \phi_{\nu-k}^{(2n-2)} \end{vmatrix}, \quad \phi_{\nu-k}^{(m)} = \left(\frac{d}{dt}\right)^m \left(e^{\frac{t^2}{4}} \varphi_{\nu-k}\right). \tag{4.43}$$

Therefore, we obtain the following proposition.

Proposition 4.3 *We define the functions $\tau_n^{\nu-k}$ by (4.43). Then,*

$$q = -\frac{\tau_n^{\nu} \tau_{n+1}^{\nu-1}}{\tau_n^{\nu-1} \tau_{n+1}^{\nu}}, \quad p = (\nu+1) \frac{\tau_n^{\nu-1} \tau_{n+1}^{\nu+1}}{\tau_{n+1}^{\nu} \tau_n^{\nu}}, \tag{4.44}$$

$$\kappa_0 = -\nu + n, \quad \theta_\infty = -\nu - 1, \tag{4.45}$$

gives a family of classical transcendental solutions of S_{IV}.

The direct derivation of this proposition is given in Appendix.

Remark 4.4 *A special case ($c_2 = 0$) of Proposition 4.3 is stated in [5]. In [6], the case of $\nu \in \mathbb{Z}$, where the τ-functions are reduced to some polynomials, is discussed.*

4.4 From $\mathrm{P_{III}}$ to $\mathrm{P_{II}}$

Both $\mathrm{P_{III}}$ and $\mathrm{P_{IV}}$ go to $\mathrm{P_{II}}$ by coalescence. First, we consider the degeneration from S_{III} to S_{II}, which is achieved by putting

$$\begin{gathered} q \to 1 + \varepsilon q, \quad p \to \varepsilon^{-1} p, \quad \theta_0 \to 2\varepsilon^{-3} + \alpha^{(1)}, \quad \theta_\infty \to -2\varepsilon^{-3} + \alpha^{(2)}, \\ t \to -2\varepsilon^{-3} \left(1 - \frac{1}{2}\varepsilon^2 t\right), \quad H \to -2\varepsilon^{-2} H - 2\varepsilon^{-3} \alpha, \end{gathered} \tag{4.46}$$

and taking the limit as $\varepsilon \to 0$, where we set $\alpha = \frac{1}{2}\left(\alpha^{(1)} + \alpha^{(2)}\right)$.

Let us consider the degeneration of the family of classical transcendental solutions. From (4.46), it is the case of $\sigma = 1$ (the Bessel function) that we can take the degeneration limit. By the relation

$$J_{-\nu} = \cos(\nu\pi) J_\nu - \sin(\nu\pi) Y_\nu, \tag{4.47}$$

we rewrite $\varphi_{\nu+j}$ (4.21) as

$$\varphi_{\nu+j} = c_1 J_{\nu+j} + c_2 Y_{\nu+j}. \tag{4.48}$$

Then, from (4.46), we see that by putting

$$\nu = -2\varepsilon^{-3}, \tag{4.49}$$

$\mathrm{S_{III}}$ with (4.25) is reduced to $\mathrm{S_{II}}$ with $\alpha = n$. It is known that we have [1]

$$\begin{aligned} J_\nu(\nu + z\nu^{\frac{1}{3}}) &= 2^{\frac{1}{3}}\nu^{-\frac{1}{3}}\mathrm{Ai}(-2^{\frac{1}{3}}z) + O(\nu^{-1}), \\ Y_\nu(\nu + z\nu^{\frac{1}{3}}) &= -2^{\frac{1}{3}}\nu^{-\frac{1}{3}}\mathrm{Bi}(-2^{\frac{1}{3}}z) + O(\nu^{-1}), \end{aligned} \tag{4.50}$$

which lead us to

$$J_\nu(t) \to -\varepsilon \mathrm{Ai}(t) + O(\varepsilon^3), \quad Y_\nu(t) \to \varepsilon \mathrm{Bi}(t) + O(\varepsilon^3). \tag{4.51}$$

Thus, we get

$$\tau_n^{\nu+j} \to \varepsilon^n (-2\varepsilon^{-2})^{(n-1)n} \tau_n, \tag{4.52}$$

where the functions τ_n are defined by (1.13). Therefore, we obtain Proposition 1.1.

4.5 From $\mathrm{P_{IV}}$ to $\mathrm{P_{II}}$

It is well known that the Hamiltonian system $\mathrm{S_{II}}$ is also derived from $\mathrm{S_{IV}}$ by degeneration. This process is achieved by putting

$$\begin{gathered} q \to \varepsilon^{-3}(1 - \varepsilon^2 q), \quad p \to -\varepsilon p, \quad t \to -2\varepsilon^{-3}\left(1 + \frac{1}{2}\varepsilon^4 t\right), \\ \kappa_0 \to \varepsilon^{-6}, \quad \theta_\infty \to \alpha, \quad H \to -\varepsilon^{-1} H + \varepsilon^{-3}\alpha, \end{gathered} \tag{4.53}$$

and taking the limit as $\varepsilon \to 0$.

Let us consider the degeneration of the classical transcendental solutions. Applying the Bäcklund transformation π to the solutions in Theorem A.2, we obtain the following corollary.

Corollary 4.5 *We define the functions τ_n^ν by (4.43). Then,*

$$q = \frac{d}{dt}\log\frac{\tau_{n+1}^\nu}{\tau_n^\nu} - t, \quad p = \frac{\tau_{n+1}^\nu \tau_{n-1}^{\nu-1}}{\tau_n^{\nu-1}\tau_n^\nu}, \tag{4.54}$$

$$\kappa_0 = \nu + 1, \quad \theta_\infty = n, \tag{4.55}$$

gives a family of classical transcendental solutions of S_{IV}.

It is easy to see that by putting

$$\nu \to \varepsilon^{-6} - 1, \tag{4.56}$$

$\mathrm{S_{IV}}$ with (4.55) reduced to $\mathrm{S_{II}}$ with $\alpha = n$ in the limit as $\varepsilon \to 0$.

Let us consider the degeneration of classical transcendental solutions. According to [3], we find that the parabolic cylinder function is reduced to the Airy function as

$$\frac{D_{\nu+j}(t)}{\Gamma(\nu+j+1)} \to (-\varepsilon)^{3j}\mathrm{Ai}(t), \quad \Gamma(-\nu-j)D_{\nu+j}(-t) \to (-\varepsilon)^{3j}\mathrm{Ai}(\omega t), \tag{4.57}$$

with $\omega = e^{2\pi i/3}$. Thus, we have

$$\varphi_{\nu+j} \to (-\varepsilon)^{3j}\varphi, \tag{4.58}$$

where φ denote the general solution of Airy's differential equation. Normalizing the τ-functions (4.43) as

$$\tau_n^{\nu+j} = e^{n\frac{t^2}{4}}\tilde{\tau}_n^{\nu+j}, \tag{4.59}$$

we get

$$\tilde{\tau}_n^{\nu+j} \to (-\varepsilon)^{-n(n-1)+3jn}\tau_n, \tag{4.60}$$

where the functions τ_n are defined by (1.13). Therefore, we obtain Proposition 1.1.

Acknowledgment The author would like to thank Prof. Y. Haraoka, Prof. K. Kajiwara and Prof. M. Noumi for their helpful comments.

A Classical transcendental solutions of the Painlevé IV equation

A.1 The symmetric form of the Painlevé IV equation

The symmetric form of $\mathrm{P_{IV}}$ is given by [6]

$$\begin{aligned} f_0' &= f_0(f_1 - f_2) + \alpha_0, \\ f_1' &= f_1(f_2 - f_0) + \alpha_1, \quad {}' = \frac{d}{dt}, \\ f_2' &= f_2(f_0 - f_1) + \alpha_2, \end{aligned} \tag{A.1}$$

with the normalization conditions $\alpha_0 + \alpha_1 + \alpha_2 = 1$ and $f_0 + f_1 + f_2 = t$. The correspondence to the canonical variables and parameters of $\mathrm{S_{IV}}$ is given by

$$q = -f_1, \quad p = f_2, \qquad \kappa_0 = \alpha_1, \quad \theta_\infty = -\alpha_2. \tag{A.2}$$

The Bäcklund transformations of $\mathrm{P_{IV}}$ are described as follows,

$$\begin{aligned} &s_i(\alpha_i) = -\alpha_i, \quad s_i(\alpha_j) = \alpha_j + \alpha_i \ \ (j = i \pm 1), \quad \pi(\alpha_j) = \alpha_{j+1}, \\ &s_i(f_i) = f_i, \qquad s_i(f_j) = f_j \pm \frac{\alpha_i}{f_i} \ \ (j = i \pm 1), \quad \pi(f_j) = f_{j+1}, \end{aligned} \tag{A.3}$$

where the subscripts $i = 0, 1, 2$ are understood as elements of $\mathbb{Z}/3\mathbb{Z}$. The Hamiltonians h_i of the system (A.1) are given by

$$h_0 = f_0 f_1 f_2 + \frac{\alpha_1 - \alpha_2}{3} f_0 + \frac{\alpha_1 + 2\alpha_2}{3} f_1 - \frac{2\alpha_1 + \alpha_2}{3} f_2, \tag{A.4}$$

and $h_i = \pi^i(h_0)$. Introducing the τ-functions τ_i as $h_i = (\log \tau_i)'$, we find that the Bäcklund transformations for the τ-functions are described by

$$s_i(\tau_j) = \tau_j \ \ (i \neq j), \quad s_i(\tau_i) = f_i \frac{\tau_{i-1}\tau_{i+1}}{\tau_i}, \quad \pi(\tau_i) = \tau_{i+1}. \tag{A.5}$$

The f-variables are recovered from the τ-functions by

$$f_i = \frac{\tau_i s_i(\tau_i)}{\tau_{i-1}\tau_{i+1}} = \frac{d}{dt}\log\frac{\tau_{i-1}}{\tau_{i+1}} + \frac{t}{3}. \tag{A.6}$$

Let us define the translation operators T_i by

$$T_1 = \pi s_2 s_1, \quad T_2 = s_1 \pi s_2, \quad T_0 = s_2 s_1 \pi, \tag{A.7}$$

which commute with each other and act on the parameters α_i as

$$T_i(\alpha_{i-1}) = \alpha_{i-1} + 1, \quad T_i(\alpha_i) = \alpha_i - 1, \quad T_i(\alpha_j) = \alpha_j \ (j \neq i-1, i). \tag{A.8}$$

Noting that $T_1 T_2 T_0 = 1$, we set

$$\tau_{k,l} = T_1^k T_2^l(\tau_0), \quad k, l \in \mathbb{Z}. \tag{A.9}$$

Then, we have from (A.6)

$$\begin{aligned} T_1^k T_2^l(f_0) &= \frac{\tau_{k,l}\tau_{k+2,l+1}}{\tau_{k+1,l+1}\tau_{k+1,l}} = \frac{d}{dt}\log\frac{\tau_{k+1,l+1}}{\tau_{k+1,l}} + \frac{t}{3}, \\ T_1^k T_2^l(f_1) &= \frac{\tau_{k+1,l}\tau_{k,l+1}}{\tau_{k,l}\tau_{k+1,l+1}} = \frac{d}{dt}\log\frac{\tau_{k,l}}{\tau_{k+1,l+1}} + \frac{t}{3}, \\ T_1^k T_2^l(f_2) &= \frac{\tau_{k+1,l+1}\tau_{k,l-1}}{\tau_{k+1,l}\tau_{k,l}} = \frac{d}{dt}\log\frac{\tau_{k+1,l}}{\tau_{k,l}} + \frac{t}{3}. \end{aligned} \tag{A.10}$$

It is possible to derive the Toda equation with respect to each translation operator. For the T_1-direction, we have

$$\tau_{k+1,l}\tau_{k-1,l} = \left[(\log \tau_{k,l})'' + \frac{2\alpha_1 + \alpha_2 - 2k + l}{3}\right]\tau_{k,l} \cdot \tau_{k,l}. \tag{A.11}$$

A.2 A Riccati solution

We derive a Riccati solution of (A.1). First, we set $\alpha_0 = 0$ and $f_0 = 0$. Then, f_1 satisfies a Riccati equation

$$f_1' = f_1(t - f_1) + \alpha_1. \tag{A.12}$$

By the dependent variable transformation $f_1 = (\log \varphi)' + \frac{t}{2}$, we have for φ the linear equation

$$\left(\frac{d^2}{dt^2} - \alpha_1 + \frac{1}{2} - \frac{t^2}{4}\right)\varphi = 0, \tag{A.13}$$

which is nothing but Weber's differential equation. We set $\alpha_1 = -\nu$ ($\notin \mathbb{Z}$). Then, the general solution of (A.13) is expressed by

$$\varphi = c_1 \frac{D_\nu(t)}{\Gamma(\nu+1)} + c_2 \Gamma(-\nu) D_\nu(-t), \tag{A.14}$$

where c_i ($i = 1, 2$) are arbitrary complex constants. By using the contiguity relations of the hyperbolic cylinder function, we obtain the following proposition.

Proposition A.1 *We define the function* $\varphi_{\nu-k}$ *by (4.41). Then,*

$$(f_0, f_1, f_2) = \left(0, \frac{\varphi_{\nu-1}}{\varphi_\nu}, (\nu+1)\frac{\varphi_{\nu+1}}{\varphi_\nu}\right), \quad (\alpha_0, \alpha_1, \alpha_2) = (0, -\nu, \nu+1), \tag{A.15}$$

gives a Riccati solution of the symmetric form of P_{IV}.

A.3 A Determinant formula for a family of classical transcendental solutions

First, we calculate the Hamiltonians and τ-functions for the Riccati solution in Proposition A.1. Under the specialization (A.15), the Hamiltonians and τ-functions are calculated as

$$h_0 = \frac{\varphi_\nu'}{\varphi_\nu} + \frac{\nu + \frac{1}{2}}{3}t, \quad h_1 = \frac{\nu+1}{3}t, \quad h_2 = \frac{\nu}{3}t, \tag{A.16}$$

and

$$\begin{aligned}
&\tau_0 = \tau_{0,0} = \varphi_\nu \exp\left(\frac{\nu+\frac{1}{2}}{6}t^2\right), \quad \tau_1 = \tau_{1,0} = \exp\left(\frac{\nu+1}{6}t^2\right), \quad \tau_2 = \tau_{1,1} = \exp\left(\frac{\nu}{6}t^2\right), \\
&s_0(\tau_0) = \tau_{2,1} = 0, \\
&s_1(\tau_1) = \tau_{0,1} = \varphi_{\nu-1}\exp\left(\frac{\nu-\frac{1}{2}}{6}t^2\right), \quad s_2(\tau_2) = \tau_{0,-1} = (\nu+1)\varphi_{\nu+1}\exp\left(\frac{\nu+\frac{3}{2}}{6}t^2\right),
\end{aligned} \tag{A.17}$$

up to multiplication by some constants, respectively. Introducing the functions $\sigma_{k,l}$ by

$$\tau_{k,l} = \sigma_{k,l}\exp\left(\frac{\nu+2k-l-1}{6}t^2\right), \tag{A.18}$$

we see that

$$\sigma_{2,l} = 0, \quad \sigma_{1,l} = \text{const.}, \quad \sigma_{0,l} = (\text{const.}) \times e^{\frac{t^2}{4}}\varphi_{\nu-l}. \tag{A.19}$$

Moreover, we set

$$\sigma_{k,l} = \omega_{k,l}\rho_{k,l}, \quad \omega_{k,l} = \omega_{k,l}(\nu), \tag{A.20}$$

with $\rho_{1,l} = 1$ and $\rho_{0,l} = e^{\frac{t^2}{4}}\varphi_{\nu-l}$, and impose that the constants $\omega_{k,l}$ satisfy

$$\omega_{k+1,l}\omega_{k-1,l} = \omega_{k,l}^2. \tag{A.21}$$

From the Toda equation (A.11), the function $\rho_{k,l}$ are determined by

$$\rho_{k+1,l}\rho_{k-1,l} = \rho_{k,l}''\rho_{k,l} - (\rho_{k,l}')^2, \tag{A.22}$$

with the initial conditions

$$\rho_{2,l} = 0, \quad \rho_{1,l} = 1, \quad \rho_{0,l} = e^{\frac{t^2}{4}}\varphi_{\nu-l}. \tag{A.23}$$

By Darboux's formula, the functions $\rho_{1-n,l}$ for $n \in \mathbb{Z}_{\geq 0}$ are expressed as

$$\rho_{1-n,l} = \begin{vmatrix} \rho_{0,l}^{(0)} & \rho_{0,l}^{(1)} & \cdots & \rho_{0,l}^{(n-1)} \\ \rho_{0,l}^{(1)} & \rho_{0,l}^{(2)} & \cdots & \rho_{0,l}^{(n)} \\ \vdots & \vdots & \ddots & \vdots \\ \rho_{0,l}^{(n-1)} & \rho_{0,l}^{(n)} & \cdots & \rho_{0,l}^{(2n-2)} \end{vmatrix}, \quad \rho_{0,l}^{(i)} = \left(\frac{d}{dt}\right)^i \rho_{0,l}. \tag{A.24}$$

Note that the constants $\omega_{k,l}$ are determined by the recurrence relations (A.21) and

$$\omega_{i,l+1}\omega_{i,l-1} = (\nu+1-l)\omega_{i,l}^2, \tag{A.25}$$

with the initial conditions $\omega_{i,0} = \omega_{i,1} = 1$ $(i = 0,1)$. Since it is possible to set $l = 0$ without loss of generality, we obtain the following theorem.

Theorem A.2 *We define the functions* $\tau_n^{\nu-k}$ *by (4.43). Then,*

$$f_0=\frac{\tau_{n+1}^{\nu}\tau_{n-1}^{\nu-1}}{\tau_n^{\nu-1}\tau_n^{\nu}}=\frac{d}{dt}\log\frac{\tau_n^{\nu-1}}{\tau_n^{\nu}},$$

$$f_1=\frac{\tau_n^{\nu}\tau_{n+1}^{\nu-1}}{\tau_{n+1}^{\nu}\tau_n^{\nu-1}}=\frac{d}{dt}\log\frac{\tau_{n+1}^{\nu}}{\tau_n^{\nu-1}}, \tag{A.26}$$

$$f_2=(\nu+1)\frac{\tau_n^{\nu-1}\tau_{n+1}^{\nu+1}}{\tau_n^{\nu}\tau_{n+1}^{\nu}}=\frac{d}{dt}\log\frac{\tau_n^{\nu}}{\tau_{n+1}^{\nu}}+t,$$

$$(\alpha_0,\alpha_1,\alpha_2)=(-n,-\nu+n,\nu+1)\,, \tag{A.27}$$

and

$$q=-\frac{\tau_n^{\nu}\tau_{n+1}^{\nu-1}}{\tau_{n+1}^{\nu}\tau_n^{\nu-1}}=-\frac{d}{dt}\log\frac{\tau_{n+1}^{\nu}}{\tau_n^{\nu-1}},\quad p=(\nu+1)\frac{\tau_n^{\nu-1}\tau_{n+1}^{\nu+1}}{\tau_n^{\nu}\tau_{n+1}^{\nu}}=\frac{d}{dt}\log\frac{\tau_n^{\nu}}{\tau_{n+1}^{\nu}}+t, \tag{A.28}$$

$$\kappa_0=-\nu+n,\quad \theta_\infty=-\nu-1, \tag{A.29}$$

gives a family of classical transcendental solutions of the symmetric form of P_{IV} *and the Hamiltonian system* S_{IV}*, respectively.*

References

[1] M. Abramowitz and I. A. Stegun, Handbook of Mathematical Functions, Dover Publications, New York, 1964.

[2] K. Iwasaki, H. Kimura, S. Shimomura and M. Yoshida, From Gauss to Painlevé – A Modern Theory of Special Functions, Aspects of Mathematics E16, Vieweg, 1991.

[3] Y. Haraoka, Confluence of cycles for hypergeometric functions on $Z_{2,n+1}$, Trans. AMS. **349** (1997) 675-712.

[4] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, Determinant formulas for the Toda and discrete Toda equations, Funkcial. Ekvac. **44** (2001) 291-307.

[5] K. Kajiwara and Y. Ohta, Determinant structure of the rational solutions for the Painlevé IV equation, J. Phys. A: Math. Gen. **31** (1998) 2431-2446.

[6] M. Noumi and Y. Yamada, Symmetries in the fourth Painlevé equation and Okamoto polynomials, Nagoya Math. J. **153** (1999) 53-86.

[7] M. Noumi and Y. Yamada, Higher order Painlevé equations of type $A_l^{(1)}$, Funkcial. Ekvac. **41** (1998) 483-503.

[8] M. Noumi and Y. Yamada, Affine Weyl groups, discrete dynamical systems and Painlevé equations, Commun. Math. Phys. **199** (1998) 281-295.

[9] M. Noumi and Y. Yamada, A new Lax pair for the sixth Painlevé equation associated with $\widehat{\mathfrak{so}}(8)$, In: Microlocal Analysis and Complex Fourier Analysis (Eds. T. Kawai and K. Fujita), World Scientific, 2002, pp. 238-252.

[10] K. Okamoto, Studies on the Painlevé equations I, sixth Painlevé equation $\mathrm{P_{VI}}$, Annali di Matematica pura ed applicata **CXLVI** (1987) 337-381.

[11] K. Okamoto, Studies on the Painlevé equations II, fifth Painlevé equation $\mathrm{P_V}$, Japan J. Math. **13** (1987) 47-76.

[12] K. Okamoto, Studies on the Painlevé equations III, second and fourth Painlevé equations, $\mathrm{P_{II}}$ and $\mathrm{P_{IV}}$, Math. Ann. **275** (1986) 222-254.

[13] K. Okamoto, Studies on the Painlevé equations IV, third Painlevé equation $\mathrm{P_{III}}$, Funkcial. Ekvac. **30** (1987) 305-332.

[14] P. Painlevé, Sur les équations différentielles du second ordre à points critiques fixes, C. R. Acad. Sci. Paris **143** (1906) 1111-1117.